

caused by pannus ingrowth and possibly caused by rupture of the support, can occur and is thus worth reporting.

*Giovanni Battista Luciani, MD
Paolo Bertolini, MD
Alessandro Mazzucco, MD
Division of Cardiac Surgery
University of Verona
Verona, Italy*

REFERENCES

1. Luciani GB, Bertolini P, Mazzucco A. Early failure of free-hand aortic stentless xenograft valves. *J Thorac Cardiovasc Surg* 1997;113:1109-10.
2. Luciani GB, Bertolini P, Vecchi B, Mazzucco A. Mid-term results after aortic valve replacement with freehand stentless xenografts: a comparison of three prostheses. *J Thorac Cardiovasc Surg*. In press.
3. Hvass U, Chatel D, Ouroudji M, et al. The O'Brien-Angell stentless valve: early results of 100 implants. *Eur J Cardiothorac Surg* 1994;8:384-7.
4. Hvass U, Chatel D, Assayag P, et al. The O'Brien-Angell stentless porcine valve: early results with 150 implants. *Ann Thorac Surg* 1995;60:S414-7.
5. O'Brien MF. The CryoLife-O'Brien composite aortic stentless xenograft: surgical technique of implantation. *Ann Thorac Surg* 1995;60:S410-3.

12/8/88051

Minimally invasive coronary artery surgery

To the Editor:

We read with great interest the editorial by Reardon and associates¹ regarding minimally invasive coronary artery surgery. We share their concerns about the safety of minimally invasive coronary artery bypass, particularly when the left internal thoracic artery is used to bypass the left anterior descending artery, our "gold standard." The operation with the best track record may have been converted into an extremely dangerous surgical tool.

Many surgeons attended meetings dealing with minimally invasive coronary artery surgery and returned with great enthusiasm. From the outset, however, we were very concerned that small thoracic incisions were not the ideal approach from the technical and safety standpoints in patients who could become unstable, regardless of whether the surgeon had access to the femoral vessels.

We have adopted the surgical exposure described by Arom, Emery, and Nicoloff.² We modified their techniques to use a complete sternotomy approach via a small skin incision, usually 5 inches (12.5 cm) in length. Because of the skin's great elasticity, we can perform a complete median sternotomy using the standard sternal saw. In this way, we have complete access to the heart via a small skin incision that is cosmetically appealing.

Through this approach we can mobilize both internal thoracic arteries and use the radial artery as a free graft to reach diagonal branches, the ramus marginalis, or proximal circumflex branches. Because we use cardiopulmonary bypass and the standard antegrade/retrograde cardioplegia with the heart still, we can construct perfect

anastomoses. So much has been written about the deleterious effects of cardiopulmonary bypass that the safety of a 5- to 15-minute pump run with cardioplegic arrest has been forgotten.

We have used this strategy and have not had a single instance of symptoms or signs of coronary insufficiency from the operations performed. We wonder whether the surgeons who have commercial interests in the companies promoting minimally invasive surgery can claim this completeness of revascularization and excellence of results. We congratulate Dr. Reardon and his group for putting a word of caution to the enthusiasm that minimally invasive coronary surgery has generated.

*Raúl García Rinaldi, MD
Ernesto R. Soltero, MD
Jorge Carballido, MD
Pavía Heart Institute
San Juan, Puerto Rico*

REFERENCES

1. Reardon MJ, Espada R, Letsou GV, et al. Editorial: minimally invasive coronary artery surgery—a word of caution. *J Thorac Cardiovasc Surg* 1997;14:419-20.
2. Arom KV, Emery RW, Nicoloff DM. Mini-sternotomy for coronary artery bypass grafting. *Ann Thorac Surg* 1996;61:1271-2.

12/8/87794

Right heart bypass with an extracardiac conduit: A cautionary tale!

To the Editor:

In this Journal, we¹ have recently commented on our use of total right heart bypass with an extracardiac conduit.

In performing this procedure, taking a generous cuff of the inferior part of the right atrial wall at its junction with the inferior caval vein, we found it possible to "upsized" the diameter of the conduit over and above the extant diameter of the caval vein at the level of the diaphragm.

A more recent experience with this surgical approach in a 4-year-old child with complex cyanotic congenital heart disease revealed a hidden trap: Rapid accumulation of ascites in the early postoperative period was accompanied by a mean gradient of 10 mm Hg between the inferior caval vein (17 mm Hg) and the conduit (7 mm Hg). The difference in pressures had been noticed at the time of the operation but, because of the external appearance of a wide trumpet-shaped lower anastomosis, the measurements had been discounted as "artifact." Because of the increasing ascites, an exploration became necessary when the gradient across the lower anastomosis was also demonstrated by direct manometry.

On takedown, we observed that a large and prominent eustachian valve had been caught up in the anastomosis, producing a partial curtain across the venous pathway. After excision of the valve and reconstruction of the anastomosis, the gradient was abolished.

When writing on prominence of the eustachian and the thebesian valves, Trento and colleagues² commented that the structures might be of “functional significance only if it were necessary to perform a Fontan procedure when they might obstruct flow through an atriopulmonary (or atrioventricular) anastomosis.” A cautionary tale!

*Christopher Lincoln, FRCS
Consultant Cardiothoracic Surgeon
Royal Brompton Hospital
London SW3 6NP, United Kingdom*

REFERENCES

1. Roberts CS, Peters P, Lincoln C. Right heart bypass with an extracardiac conduit for functionally single ventricle with anomalous pulmonary venous connection to the right atrium. *J Thorac Cardiovasc Surg* 1997;114:297-8.
2. Trento A, Zuberbuhler JR, Anderson RH, Park SC, Siewers RD. Divided right atrium (prominence of the eustachian and thebesian valves). *J Thorac Cardiovasc Surg* 1988;96:457-63.

12/8/88049